

Agroforestry systems: Meta-analysis of soil carbon stocks, sequestration processes, and future potentials

Shi L., Feng W., Xu J., Kuzyakov Y.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 John Wiley & Sons, Ltd. Agroforestry (AF) has the potential to restore degraded lands, provide a broader range of ecosystem goods and services such as carbon (C) sequestration and high biodiversity, and increase soil fertility and ecosystem stability through additional C input from trees, erosion prevention, and microclimate improvement. Advantages and processes for global C sequestration in AF are unknown. We used a meta-analysis of 427 soil C stock data pairs grouped into four main AF systems—alley cropping, windbreaks, silvopastures, and homegardens—and evaluated changes in AF and adjacent control cropland or pasture. Mean soil C stocks in AF (1-m depth) were 126 Mg C·ha⁻¹, which is 19% more than that in cropland or pasture. The highest C stocks in soil were in subtropical homegardens, AF with younger trees, and topsoil (0–20 cm). Increased soil C stocks in AF were lower than aboveground C stocks in most AF systems, except alley cropping. Homegardens stored the highest C in both aboveground and belowground, especially in the subsoil (20–100 cm). Advantages of AF ecosystem services focusing on mechanisms of belowground C sequestration were analyzed. AF could store 5.3×10^9 Mg additional C in soil on 944 Mha globally, with most in the tropics and subtropics. AF systems could greatly contribute to global soil C sequestration if used in larger areas. Future investigations of AF should include (a) mechanistic- and process-based studies (instead of common monitoring and inventories), (b) models linking forest and crop growth with soil water and C and nutrient cycling, and (c) accurate assessments of the AF area worldwide based on the remote sensing approaches.

<http://dx.doi.org/10.1002/ldr.3136>

Keywords

agroforestry management, carbon sequestration, ecosystem services, homegardens, meta-analysis, sustainable land use

References

- [1] Albrecht, A., & Kandji, S. T. (2003). Carbon sequestration in tropical agroforestry systems. *Agriculture Ecosystems and Environment*, 99, 15–27. [https://doi.org/10.1016/S0167-8809\(03\)00138-5](https://doi.org/10.1016/S0167-8809(03)00138-5)
- [2] Araujo, A. S. F., Iwata, B. D. F., de Andrade Lira, M. Jr., & Xavier, G. R. (2012). Microbiological process in agroforestry systems. A review. *Agronomy for Sustainable Development*, 32, 215–226. <https://doi.org/10.1007/s13593-011-0026-0>
- [3] Arora, G., Chaturvedi, S., Kaushal, R., Nain, A., Tewari, S., Alam, N. M., & Chaturvedi, O. P. (2014). Growth, biomass, carbon stocks, and sequestration in age series of *Populus deltoides* plantations in Tarai region of central Himalaya. *Turkish Journal of Agriculture and Forestry*, 38, 550–560. <https://doi.org/10.3906/tar-1307-94>

- [4] Atangana A, Khasa D, Chang S, Ann Degrande 2014. Agroforestry for soil conservation. Tropical agroforestry. Dordrecht: Springer. 2014: 203-216
- [5] Baah-Acheamfour, M., Chang, S. X., Bork, E. W., & Carlyle, C. N. (2017). The potential of agroforestry to reduce atmospheric greenhouse gases in Canada: Insight from pairwise comparisons with traditional agriculture, data gaps and future research. *Forestry Chronicle*, 93, 180-189. <https://doi.org/10.5558/tfc2017-024>
- [6] Balakrishna, A. N., Lakshmipathy, R., Bagyaraj, D. J., & Ashwin, R. (2017). Influence of alley cropping system on am fungi, microbial biomass C and yield of finger millet, peanut and pigeon pea. *Agroforestry Systems*, 91, 487-493. <https://doi.org/10.1007/s10457-016-9949-4>
- [7] Basu, J. P. (2014). Agroforestry, climate change mitigation and livelihood security in India. *New Zealand Journal of Forestry Science*, 44, 1-10. <https://doi.org/10.1186/1179-5395-44-S1-S11>
- [8] Becker, J., Pabst, H., Mnyonga, J., & Kuzyakov, Y. (2015). Annual litterfall dynamics and nutrient deposition depending on elevation and land use at Mt. Kilimanjaro. *Biogeosciences*, 12, 5635-5646. <https://doi.org/10.5194/bg-12-5635-2015>
- [9] Béliveau, A., Lucotte, M., Davidson, R., Paquet, S., Mertens, F., Passos, C. J., & Romana, C. A. (2017). Reduction of soil erosion and mercury losses in agroforestry systems compared to forests and cultivated fields in the Brazilian Amazon. *Journal of Environmental Management*, 203, 522-532. <https://doi.org/10.1016/j.jenvman.2017.07.037>
- [10] Boffa, J. M., Taonda, S. J. B., Dickey, J. B., & Boffa, J. M. (2000). Field-scale influence of karite (*Vitellaria paradoxa*) on sorghum production in the Sudan zone of Burkina Faso. *Agroforestry Systems*, 49, 153-175. <https://doi.org/10.1023/a:1006389828259>
- [11] Brandle, J. R., Hodges, L., & Zhou, X. H. (2004). Windbreaks in North American agricultural systems. *New Vistas in Agroforestry*. Springer Netherlands. <https://doi.org/10.1007/978-94-017-2424-15>
- [12] Brown, D. R. (2006). Personal preferences and intensification of land use: Their impact on southern Cameroonian slash-and-burn agroforestry systems. *Agroforestry Systems*, 68, 53-67. <https://doi.org/10.1007/s10457-006-0003-9>
- [13] Cardinael, R., Chevallier, T., Cambou, A., Béral, C., Barthès, B. G., Dupraz, C., ... Kouakoua, E. C. C. (2017). Increased soil organic carbon stocks under agroforestry: A survey of six different sites in France. *Agriculture, Ecosystems and Environment*, 236, 243-255. <https://doi.org/10.1016/j.AGEE.2016.12.011>
- [14] Cardinael, R., Mao, Z., Prieto, I., Stokes, A., Dupraz, C., Kim, J. H., & Jourdan, C. (2015). Competition with winter crops induces deeper rooting of walnut trees in a Mediterranean alley cropping agroforestry system. *Plant and Soil*, 391, 219-235. <https://doi.org/10.1007/s11104-015-2422-8>
- [15] Dixon, R. K., Winjum, J. K., Andrasko, K. J., Lee, J. J., & Schroeder, P. E. (1994). Integrated land-use systems: Assessment of promising agroforest and alternative land-use practices to enhance carbon conservation and sequestration. *Climatic Change*, 27, 71-92. <https://doi.org/10.1007/bf01098474>
- [16] Gamble, J. D., Johnson, G., Sheaffer, C. C., Current, D. A., & Wyse, D. L. (2014). Establishment and early productivity of perennial biomass alley cropping systems in Minnesota, USA. *Agroforestry Systems*, 88, 75-85. <https://doi.org/10.1007/s10457-013-9657-2>
- [17] Gelaw, A. M., Singh, B. R., & Lal, R. (2015). Organic carbon and nitrogen associated with soil aggregates and particle sizes under different land uses in tigray, northern Ethiopia. *Land Degradation & Development*, 26, 690-700. <https://doi.org/10.1002/ldr.2261>
- [18] Guimaraes, G. P., Mendona, E. D., Passos, R. R., & Andrade, F. V. (2014). Soil aggregation and organic carbon of Oxisols under coffee in agroforestry systems. *Revista Brasileira De Ciencia Do Solo*, 38, 278-287. <https://doi.org/10.1590/S0100-06832014000100028>
- [19] Gunina, A., & Kuzyakov, Y. (2014). Pathways of litter C by formation of aggregates and SOM density fractions: Implications from C natural abundance. *Soil Biology and Biochemistry*, 71, 95-104. <https://doi.org/10.1590/S0100-06832014000100028>
- [20] Gurevitch, J., Curtis, P. S., & Jones, M. H. (2001). Meta-analysis in ecology. *Advances in Ecological Research*, 32, 199-247. [https://doi.org/10.1016/S0065-2504\(01\)32013-5](https://doi.org/10.1016/S0065-2504(01)32013-5)
- [21] Hemp, A. (2005). The Chagga home gardens—Relict areas for endemic Saltatoria species (Insecta: Orthoptera) on Mount Kilimanjaro. *Biological Conservation*, 125, 203-209. <https://doi.org/10.1016/j.biocon.2005.03.018>
- [22] Hemp, A. (2006). Vegetation of Kilimanjaro: Hidden endemics and missing bamboo. *African Journal of Ecology*, 44, 305-328. <https://doi.org/10.1111/j.1365-2028.2006.00679.x>
- [23] Hemp C, Hemp A. (2008). The chagga homegardens on Kilimanjaro. *Magazine of the International Human Dimensions Programme on Global Environmental Change. Mountainous Regions: Laboratories for Adaptation* 12-17. <https://doi.org/10.1016/j.biocon.2005.03.018>
- [24] Ilany, T., Ashton, M. S., Montagnini, F., & Martinez, C. (2010). Using agroforestry to improve soil fertility: effects of intercropping on *Ilex paraguariensis* (yerba mate) plantations with *Araucaria angustifolia*. *Agroforestry Systems*, 80, 399-409. <https://doi.org/10.1007/s10457-010-9317-8>

- [25] Islam, M., Dey, A., & Rahman, M. (2015). Effect of tree diversity on soil organic carbon content in the homegarden agroforestry system of north-eastern Bangladesh. *Small-Scale Forestry*, 14, 91–101. <https://doi.org/10.1007/s11842-014-9275-5>
- [26] Janssens, I. A., Dieleman, W., Luyssaert, S., Subke, J.-A., Reichstein, M., Ceulemans, R., ... Matteucci, G. (2010). Reduction of forest soil respiration in response to nitrogen deposition. *Nature Geoscience*, 3, 315–322. <https://doi.org/10.1038/ngeo844>
- [27] Kaiser, C., Koranda, M., Kitzler, B., Fuchslueger, L., Schnecker, J., Schweiger, P., ... Richter, A. (2010). Belowground carbon allocation by trees drives seasonal patterns of extracellular enzyme activities by altering microbial community composition in a beech forest soil. *New Phytologist*, 187, 843–858. <https://doi.org/10.1111/j.1469-8137.2010.03321.x>
- [28] Kaiser, K., & Kalbitz, K. (2012). Cycling downwards—Dissolved organic matter in soils. *Soil Biology and Biochemistry*, 52, 29–32. <https://doi.org/10.1016/j.soilbio.2012.04.002>
- [29] Kim, D. G., Kirschbaum, M. U. F., & Beedy, T. L. (2016). Carbon sequestration and net emissions of CH and NO under agroforestry: Synthesizing available data and suggestions for future studies. *Agriculture Ecosystems and Environment*, 226, 65–78. <https://doi.org/10.1016/j.agee.2016.04.011>
- [30] Kukal, S. S., & Bawa, S. S. (2014). Soil organic carbon stock and fractions in relation to land use and soil depth in the degraded Shiwaliks hills of lower Himalayas. *Land Degradation & Development*, 25, 407–416. <https://doi.org/10.1002/ldr.2151>
- [31] Kumar, B. M. (2011). Species richness and aboveground carbon stocks in the homegardens of central Kerala, India. *Agriculture, Ecosystems and Environment*, 140, 430–440. <https://doi.org/10.1016/j.agee.2011.01.006>
- [32] Kumar, B. M., & Nair, P. K. R. (2004). The enigma of tropical homegardens. *Agroforestry Systems*, 62, 135–152. <https://doi.org/10.1023/B:AGFO.0000028995.13227.ca>
- [33] Lacoste, M., Viaud, V., Michot, D., & Walter, C. (2015). Landscape-scale modelling of erosion processes and soil carbon dynamics under land-use and climate change in agroecosystems. *European Journal of Soil Science*, 66, 780–791. <https://doi.org/10.1111/ejss.12267>
- [34] Le Bissonnais, Y., Prieto, I., Roumet, C., Nespoulous, J., Metayer, J., Huon, S., ... Stokes, A. (2017). Soil aggregate stability in Mediterranean and tropical agro-ecosystems: Effect of plant roots and soil characteristics. *Plant and Soil*, 424, 303–317. <https://doi.org/10.1007/s11104-017-3423-6>
- [35] Lenka, N. K., Choudhury, P. R., Sudhishri, S., Dass, A., & Patnaik, U. S. (2012). Soil aggregation, carbon build up and root zone soil moisture in degraded sloping lands under selected agroforestry based rehabilitation systems in eastern India. *Agriculture Ecosystems and Environment*, 150, 54–62. <https://doi.org/10.1016/j.agee.2012.01.003>
- [36] Lorenz, K., & Lal, R. (2014). Soil organic carbon sequestration in agroforestry systems. A review. *Agronomy for Sustainable Development*, 34, 443–454. <https://doi.org/10.1007/s13593-014-0212-y>
- [37] Luo, Z., Wang, E., Feng, W., Luo, Y., & Baldock, J. (2017). The importance and requirement of belowground carbon inputs for robust estimation of soil organic carbon dynamics: Reply to Keel et al. (2017). *Global Change Biology*, 2017, 1–2. <https://doi.org/10.1111/gcb.13949>
- [38] Mann, L. K. (1986). Changes in soil carbon storage after cultivation. *Soil Science*, 142, 279–288. <https://doi.org/10.1097/00010694-198611000-00006>
- [39] Mattsson, E., Ostwald, M., Nissanka, S. P., & Marambe, B. (2013). Homegardens as a multi-functional land-use strategy in Sri Lanka with focus on carbon sequestration. *Ambio*, 42, 892–902. <https://doi.org/10.1007/s13280-013-0390-x>
- [40] Mattsson, E., Ostwald, M., Nissanka, S. P., & Pushpakumara, D. K. N. G. (2015). Quantification of carbon stock and tree diversity of homegardens in a dry zone area of Moneragala District, Sri Lanka. *Agroforestry Systems*, 89, 435–445. <https://doi.org/10.1007/s10457-014-9780-8>
- [41] Mganga, K. Z., & Kuzyakov, Y. (2014). Glucose decomposition and its incorporation into soil microbial biomass depending on land use in Mt. Kilimanjaro ecosystems. *European Journal of Soil Biology*, 62, 74–82. <https://doi.org/10.1016/j.ejsobi.2014.02.015>
- [42] Mganga, K. Z., Razavi, B. S., & Kuzyakov, Y. (2016). Land use affects soil biochemical properties in Mann LK. 1986. Changes in soil carbon storage after cultivation. *Soil Science*, 142, 279–288. <https://doi.org/10.1097/00010694-198611000-00006>
- [43] Nair, P. K. R. (1991). State-of-the-art of agroforestry systems. *Forest Ecology and Management*, 45, 5–29. [https://doi.org/10.1016/0378-1127\(91\)90203-8](https://doi.org/10.1016/0378-1127(91)90203-8)
- [44] Nair, P. K. (1993). Ramachandran. An introduction to agroforestry. Springer Science & Business Media
- [45] Nair, P. K. R. (2011). Agroforestry systems and environmental quality: Introduction. *Journal of Environmental Quality*, 40, 784–790. <https://doi.org/10.2134/jeq2011.0076>
- [46] Nair, P. K. R. (2012). Carbon sequestration studies in agroforestry systems: A reality-check. *Agroforestry Systems*, 86, 243–253. <https://doi.org/10.1007/s10457-011-9434-z>

- [47] Nair, P. K. R. (2013). Agroforestry: Trees in support of sustainable agriculture. In S. A. Elias, S. Marshall, M. Goldstein, et al. (Eds.), Reference module in Earth systems and environmental sciences. 10.1016/B978-0--2-409548-9.05088-0
- [48] Nair, P. K. R., Kumar, B. M., & Nair, V. D. (2009a). Agroforestry as a strategy for carbon sequestration. *Journal of Plant Nutrition and Soil Science*, 172, 10–23. <https://doi.org/10.1002/jpln.200800030>
- [49] Nair, P. K. R., Nair, V. D., Kumar, B. M., Haile, S. G., Dilly, O., & Pannell, D. (2009b). Soil carbon sequestration in tropical agroforestry systems: A feasibility appraisal. *Environmental Science and Policy*, 12, 1099–1111. <https://doi.org/10.1016/j.envsci.2009.01.010>
- [50] Nair, P. K. R., Saha, S. K., Nair, V. D., & Haile, S. G. (2011). Potential for greenhouse gas emissions from soil carbon stock following biofuel cultivation on degraded lands. *Land Degradation & Development*, 22, 395–409. <https://doi.org/10.1002/ldr.1016>
- [51] Nair, P. R., & Nair, V. D. (2003). Carbon storage in North American agroforestry systems. In M. J. Kimble, R. Lal, R. Birdsey, & L. S. Heath (Eds.), 2003The potential of US forest soils to sequester carbon and mitigate the greenhouse effect (pp. 333–346). New York: CRC Press LLC
- [52] Nair, V. D., & Graetz, D. A. (2004). Agroforestry as an approach to minimizing nutrient loss from heavily fertilized soils: The Florida experience. *Agroforestry Systems*, 61, 269–279. <https://doi.org/10.1023/B:AGFO.0000029004.03475.1d>
- [53] Oelbermann, M., Voroney, R. P., & Gordon, A. M. (2004). Carbon sequestration in tropical and temperate agroforestry systems: A review with examples from Costa Rica and southern Canada. *Agriculture Ecosystems and Environment*, 104, 359–377. <https://doi.org/10.1016/j.agee.2004.04.001>
- [54] Pabst, H., Gerschlauser, F., Kiese, R., & Kuzyakov, Y. (2016). Land use and precipitation affect organic and microbial carbon stocks and the specific metabolic quotient in soils of eleven ecosystems of Mt. Kilimanjaro, Tanzania. *Land Degradation & Development*, 27, 592–602. <https://doi.org/10.1002/ldr.2406>
- [55] Pandey, C. B., & Srivastava, R. C. (2009). Plant available phosphorus in homegarden and native forest soils under high rainfall in an equatorial humid tropics. *Plant and Soil*, 316, 71–80. <https://doi.org/10.1007/s11104-008-9759-1>
- [56] Pandey, D. N. (2002). Carbon sequestration in agroforestry systems. *Climate Policy*, 2, 367–377. <https://doi.org/10.3763/cpol.2002.0240>
- [57] Pausch, J., & Kuzyakov, Y. (2018). Carbon input by roots into the soil: Quantification of rhizodeposition from root to ecosystem scale. *Global Change Biology*, 24, 1–12. <https://doi.org/10.1111/gcb.13850>
- [58] Post, W. M., & Kwon, K. C. (2000). Soil carbon sequestration and land-use change: Processes and potential. *Global Change Biology*, 6, 317–327. <https://doi.org/10.1046/j.1365-2486.2000.00308.x>
- [59] Rillig, M. C. (2004). Arbuscular mycorrhizae, glomalin, and soil aggregation. *Canadian Journal of Soil Science*, 84, 355–363. <https://doi.org/10.4141/S04-003>
- [60] Saha, K. S., Nair, P. K. R., Nair, D. V., & Kumar, M. B. (2009). Soil carbon stock in relation to plant diversity of homegardens in Kerala, India. *Agroforestry Systems*, 76, 53–65. <https://doi.org/10.1007/s10457-009-9228-8>
- [61] Schmitt-Harsh, M., Evans, T. P., Castellanos, E., & Randolph, J. C. (2012). Carbon stocks in coffee agroforests and mixed dry tropical forests in the western highlands of Guatemala. *Agroforestry Systems*, 86, 141–157. <https://doi.org/10.1007/s10457-012-9549-x>
- [62] Schneckenberger, K., & Kuzyakov, Y. (2007). Carbon sequestration under *Miscanthus* in sandy and loamy soils estimated by natural C abundance. *Journal of Plant Nutrition and Soil Science*, 170, 538–542. <https://doi.org/10.1002/jpln.200625111>
- [63] Sharrow, S. H. (1999). Silvopastoralism: competition and facilitation between trees, livestock and improved grass-clover pastures on temperate rainfed lands. In: L. E. Buck, J. P. Lassoie, and E. C. M. Fernandes, (eds.) *Agroforestry in Sustainable Agricultural Systems*. USA: CRC Press, Boca Raton, FL
- [64] Singh, B., & Gill, R. I. S. (2014). Carbon sequestration and nutrient removal by some tree species in an agrisilviculture system in Punjab, India. *Range Management and Agroforestry*, 35, 107–114
- [65] Sommer, J., Dippold, M. A., Flessa, H., & Kuzyakov, Y. (2016). Allocation and dynamics of C and N within plant–soil system of ash and beech. *Journal of Plant Nutrition and Soil Science*, 179, 376–387. <https://doi.org/10.1002/jpln.201500384>
- [66] Stavi, I., & Lal, R. (2013). Agroforestry and biochar to offset climate change: A review. *Agronomy for Sustainable Development*, 33, 81–96. <https://doi.org/10.1007/s13593-012-0081-1>
- [67] Takimoto, A., Nair, P. K. R., & Nair, V. D. (2008). Carbon stock and sequestration potential of traditional and improved agroforestry systems in the West African Sahel. *Agriculture Ecosystems and Environment*, 125, 159–166. <https://doi.org/10.1016/j.agee.2007.12.010>
- [68] Takimoto, A., Nair, V. D., & Nair, P. K. R. (2009). Contribution of trees to soil carbon sequestration under agroforestry systems in the West African Sahel. *Agroforestry Systems*, 76, 11–25. <https://doi.org/10.1007/s10457-008-9179-5>

- [69] Thakur, S., Kumar, B. M., & Kunhamu, T. K. (2015). Coarse root biomass, carbon, and nutrient stock dynamics of different stem and crown classes of silver oak (*Grevillea robusta*, A. Cunn. ex R. Br.) plantation in central Kerala, India. *Agroforestry Systems*, 89, 869–883. <https://doi.org/10.1007/s10457-015-9821-y>
- [70] Torralba, M. (2016). Do European agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis. *Agriculture Ecosystems and Environment*, 230, 150–161. <https://doi.org/10.1016/j.agee.2016.06.002>
- [71] Tsonkova, P., Böhm, C., Quinkenstein, A., Freese, D., & Jose, S. (2012). Ecological benefits provided by alley cropping systems for production of woody biomass in the temperate region: a review. *Agroforestry Systems*, 85, 133–152. <https://doi.org/10.1007/s10457-012-9494-8>
- [72] Udawatta, R. P., & Jose, S. (2011). Carbon sequestration potential of agroforestry practices in temperate North America. In *Carbon sequestration potential of agroforestry systems* (pp. 17–42). Dordrecht: Springer
- [73] Upson, M. A., & Burgess, P. J. (2013). Soil organic carbon and root distribution in a temperate arable agroforestry system. *Plant and Soil*, 373, 43–58. <https://doi.org/10.1007/s11104-013-1733-x>
- [74] Upson, M. A., Burgess, P. J., & Morison, J. I. L. (2016). Soil carbon changes after establishing woodland and agroforestry trees in a grazed pasture. *Geoderma*, 283, 10–20. <https://doi.org/10.1016/j.geoderma.2016.07.002>
- [75] Vallejo, V. E., Roldan, F., & Dick, R. P. (2010). Soil enzymatic activities and microbial biomass in an integrated agroforestry chronosequence compared to monoculture and a native forest of Colombia. *Biology and Fertility of Soils*, 46, 577–587. <https://doi.org/10.1007/s00374-010-0466-8>
- [76] Verchot, N. L., Noordwijk, V. M., Kandji, S., Tomich, T., Ong, C., Albrecht, A., et al. (2007). Climate change: Linking adaptation and mitigation through agroforestry. *Mitigation and Adaptation Strategies for Global Change*, 12, 901–918. <https://doi.org/10.1007/s11027-007-9105-6>
- [77] Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36, 1–48. <https://doi.org/10.18637/jss.v036.i03>
- [78] Wiesmeier, M., Lungu, M., Cerbari, V., Boincean, B., Hübner, R., & Kögel-Knabner, I. (2018). Rebuilding soil carbon in degraded steppe soils of Eastern Europe: The importance of windbreaks and improved cropland management. *Land Degradation & Development*, 29, 875–883. <https://doi.org/10.1002/ldr.2902>
- [79] Zang, H., Blagodatskaya, E., Wen, Y., Xu, X., Dyckmans, J., & Kuzyakov, Y. (2018). Carbon sequestration and turnover in soil under the energy crop *Miscanthus*: Repeated C natural abundance approach and literature synthesis. *Global Change Biology. Bioenergy*, 10, 262–271. <https://doi.org/10.1111/gcbb.12485>
- [80] Zang, H., Wang, J., & Kuzyakov, Y. (2016). N fertilization decreases soil organic matter decomposition in the rhizosphere. *Applied Soil Ecology*, 108, 47–53. <https://doi.org/10.1016/j.apsoil.2016.07.021>
- [81] Zheng, X., Zhu, J., & Xing, Z. (2016). Assessment of the effects of shelterbelts on crop yields at the regional scale in northeast China. *Agricultural Systems*, 143, 49–60. <https://doi.org/10.1016/j.agsy.2015.12.008>
- [82] Zomer, R. J., Neufeldt, H., Xu, J. C., Ahrends, A., Bossio, D., Trabucco, A., ... Wang, M. C. (2016). Global tree cover and biomass carbon on agricultural land: The contribution of agroforestry to global and national carbon budgets. *Scientific Reports*, 6(29987). <https://doi.org/10.1038/srep29987>
- [83] Zomer RJ, Trabucco A, Coe R, Place F. (2009). *Trees on farm: Analysis of global extent and geographical patterns of agroforestry*. ICRAF Working Paper no 89. Nairobi, Kenya: World Agroforestry Centre